

DOSIMETRY IN THERAPY OF METASTATIC DIFFERENTIATED THYROID CANCER ADMINISTERING HIGH ¹³¹I ACTIVITY.

Baroli, Alberto¹; Pedrazzini, Luca¹; Lomuscio, Giuseppe²; Di Mauro, Enzo¹; Bianchi, Lorenzo⁴; Pepe, Annalisa⁴; Solbiati, Luigi⁵; Chiesa, Carlo³

¹General Hospital of Busto Arsizio, Unit of Nuclear Medicine, Busto Arsizio, Varese, Italy; ²General Hospital of Busto Arsizio, Busto Arsizio, Varese, Italy; ³Nuclear Medicine Unit, Fondazione IRCCS Istituto Nazionale Tumori Milano, Milan, Italy; ⁴General Hospital of Busto Arsizio, Unit of Health Physics, Busto Arsizio, Varese, Italy; ⁵General Hospital of Busto Arsizio, Unit of Interventional Oncologic Unit, Busto Arsizio, Varese, Italy

Background/Purpose: To evaluate the impact of ¹³¹I high activity therapy treatments of Metastatic Differentiated Thyroid Cancer (MDTC) in terms of feasibility, tolerance, efficacy, and the role that dosimetry can play

Methods: 17 patients underwent treatment with ¹³¹I high activity. Activity ranged from 6.2 GBq to 24.1 GBq. 8 of them had multiple treatments, for a total of 27 treatments.

Red marrow (RM), blood (BL) and metastases (45 in total) peri-treatment dosimetry was performed. In 12 cases, also prospective red marrow dosimetry was performed to comply the 2 Gy RM dose constraint.

Results: Only transient toxicity was observed. RM mean absorbed dose was 1.48 (0.36 - 6.67) Gy, 1.91 (0.43 - 8.71) Gy for BL, 153 (1.1 - 778) Gy for metastases. In 20/23 cases of repeated treatments, a dose reduction per unit of administered activity was observed in treatments following the first one, so a first treatment with the highest tolerable activity seems to be desirable to obtain the maximum efficacy. In 9 cases RM dose showed <50% discrepancies between prospective and peri-treatment dosimetry evaluations.

Discussion & Conclusion: Metastases from DTC became progressively less iodine-avid after repeated treatments, so a first treatment with the highest tolerable activity seems to be desirable to obtain the maximum efficacy.

In our experience, high activity treatments were well-tolerated, without persistent haematological side effects or worsening quality of life.